VOL. 458 JOURNAL OF CHROMATOGRAPHY DECEMBER 23, 1988

CONTENTS

12TH INTERNATIONAL SYMPOSIUM ON COLUMN LIQUID CHROMATOGRAPHY, WASHINGTON, DC, JUNE 19–24, 1988, PART I

Foreword
by G. Guiochon XI

Combined lectin-affinity and metal-interaction chromatography for the separation of glycophorins by high-performance liquid chromatography
by D. Corradini, Z. El Rassi, Cs. Horváth, G. Guerra and W. Horne (New Haven, CT, U.S.A.) 1

Determination of binding equilibrium constants by numerical simulation in zonal high-performance affinity chromatography
by C. Vidal-Madjar, A. Jaulmes, M. Racine and B. Sébille (Thiais, France) 13

High-performance liquid chromatography of amino acids, peptides and proteins. LXXXVII. Comparison of retention and bandwidth properties of proteins eluted by gradient and isocratic anion-exchange chromatography
by M. T. W. Hearn, A. N. Hodder and M. I. Aguilar (Clayton, Australia) 27

High-performance liquid chromatography of amino acids, peptides and proteins. LXXXVIII. Calculation of the average distance between protein solutes and the stationary phase during isocratic anion-exchange chromatography
by M. T. W. Hearn, A. N. Hodder and M. I. Aguilar (Clayton, Australia) 45

Salt-binding effects in hydrophobic-interaction chromatography
by R. A. Barford, T. F. Kumosinski, N. Parris and A. E. White (Philadelphia, PA, U.S.A.) 57

Strategy for the immobilization of monoclonal antibodies on solid-phase supports
by R. S. Matson and M. C. Little (Richmond, CA, U.S.A.) 67

Sample size and retention values in high-performance liquid chromatography of biological and synthetic polymers
by H. Engelhardt, M. Czok, R. Schultz and E. Schweinheim (Saarbrücken, F.R.G.) . . . 79

Thermodynamics of α-lactalbumin denaturation in hydrophobic-interaction chromatography and stationary phases comparison
by K. Benedek (King of Prussia, PA, U.S.A.) 93

New high-performance liquid chromatography-based methodology for monitoring the conformational transitions of self-associating hydrophobic peptides, incorporated into liposomes
by M. C. Bañó, L. Braco and C. Abad (Valencia, Spain) 105

High-performance liquid chromatographic analysis of DNA composition and DNA modification of chloroacetaldehyde
by R. P. Singhal and J. P. Landes (Wichita, KS, U.S.A.) 117

High-performance liquid chromatographic separation of peptides on a diol-Gly-Phe-Phe tripeptide-bonded phase
by T. C. Pinkerton and K. A. Koeplinger (Kalamazoo, MI, U.S.A.) 129

Computer simulation of high-performance liquid chromatographic separations of peptide and protein digests for development of size-exclusion, ion-exchange and reversed-phase chromatographic methods
by R. S. Hodges, J. M. R. Parker, C. T. Mant and R. R. Sharma (Edmonton, Canada) . 147

Protein immobilization on silica supports. A ligand density study
by D. Wu (Ames, IA, U.S.A.) and R. R. Walters (Kalamazoo, MI, U.S.A.) 169

Automated high-performance liquid chromatographic method for the determination of iodotyrosines and iodothyronines
by P. R. Kootstra (Amsterdam, The Netherlands), H. H. van den Broek, E. A. Hogendoorn and C. E. Goewie (Bilthoven, The Netherlands) and J. J. M. de Vijlder (Amsterdam, The Netherlands) 175

High-performance affinity isolation of lymphocyte membrane receptors on biotinylated antigen and avidin-coated beads
by T. M. Phillips and S. C. Frantz (Washington, DC, U.S.A.), J. V. Babashak (Vineland, NJ, U.S.A.) and J. J. Chmielinska (Washington, DC, U.S.A.) 185

Effect of peptide chain length on peptide retention behaviour in reversed-phase chromatography
by C. T. Mant, T. W. L. Burke, J. A. Black and R. S. Hodges (Edmonton, Canada) . . . 193

Rapid high-performance affinity chromatography on micropellicular sorbents
by L. Várady, K. Kalghatgi and Cs. Horváth (New Haven, CT, U.S.A.) 207

Determination of pterins in urine by high-performance liquid chromatography on C_{18} columns, conditioned with cetyl-trimethyl ammonium bromide
by B. C. Montalvo, C. I. Villar, R. C. I. Hornillos and L. P. Diez (Madrid, Spain) . . . 217

Stabilization of reversed phases for liquid chromatography. Applications of infrared spectroscopy for the study of bonded-phase stability
by N. Sagliano, Jr. and R. A. Hartwick (Piscataway, NJ, U.S.A.) and R. E. Patterson, B. A. Woods, J. L. Bass and N. T. Miller (Lafayette Hill, PA, U.S.A.) 225

Anion-exchange selectivity in latex-based columns for ion chromatography
by R. W. Slingsby and C. A. Pohl (Sunnyvale, CA, U.S.A.) 241

Electron spin resonance studies under dynamic mobile phase conditions on chemically modified silica
by C. Miller, R. Dadoo, R. G. Kooser and J. Gorse (Galesburg, IL, U.S.A.) 255

Insights into the slurry packing and bed structure of capillary liquid chromatographic columns
by D. C. Shelly, V. L. Antonucci, T. J. Edkins and T. J. Dalton (Hoboken, NJ, U.S.A.) . 267

New packing and column for fast protein high-performance liquid chromatography
by T. J. Szczerba, D. N. Baehr, L. J. Glunz and J. A. Perry (Morton Grove, IL, U.S.A.) and M. J. Holdoway (Hawarden, U.K.) 281

Liquid chromatographic determination of low-molecular-weight amides in pharmaceutical matrices
by S. V. Snorek, B. A. Olsen and D. A. Pierson (Lafayette, IN, U.S.A.) 287

High-performance liquid chromatography of the antitumour agent triethylenethiophosphoramide and its metabolite triethylenephopsphoramide with sodium sulphide, taurine and *o*-phthalaldehyde as pre-column fluorescent derivatization reagents
by A. Sano, S. Matsutani and S. Takitani (Tokyo, Japan) 295

Separation and detection of DNA by capillary electrophoresis
by T. J. Kasper, M. Melera, P. Gozel and R. G. Brownlee (Sunnyvale, CA, U.S.A.) . . . 303

On-line capillary zone electrophoresis–ion spray tandem mass spectrometry for the determination of dynorphins
E. D. Lee, W. Mück and J. D. Henion (Ithaca, NY, U.S.A.) and T. R. Covey (Thornhill, Canada) 313

Rapid separation of DNA restriction fragments using capillary electrophoresis
by A. S. Cohen, D. Najarian, J. A. Smith and B. L. Karger (Boston, MA, U.S.A.) . . . 323

Strategies for automated optimisation of high-performance liquid chromatographic separations incorporating diode-array detection
by A. G. Wright and A. F. Fell (Bradford, U.K.) and J. C. Berridge (Kent, U.K.) 335

Correction of the resolution function for non-ideal peaks
by P. J. Schoenmakers (Eindhoven, The Netherlands) and J. K. Strasters and Á. Bartha (Delft, The Netherlands) 355

Practical optimization of the separation of a limited subset of components by using isoeluotropic ternary eluent mixtures in reversed-phase high-performance liquid chromatography
by Á. Bartha, H. A. H. Billiet and L. de Galan (Delft, The Netherlands) 371

Computer-assisted retention prediction system for oligonucleotides in gradient anion-exchange chromatography
by Y. Baba and M. Fukuda (Oita, Japan) and N. Yoza (Fukuoka, Japan) 385

Determination of hydrophobicity parameters on polybutadiene-coated alumina and their application in quantitative structure–activity relationships analysis
by R. Kaliszan and J. Petrusewicz (Gdansk, Poland) and R. W. Blain and R. A. Hartwick (Piscataway, NJ, U.S.A.) 395

Author Index 405

In articles with more than one author, the name of the author to whom correspondence should be addressed is indicated in the article heading by a 6-pointed asterisk (*)